

# **Epiomic Epidemiology Series: Dysmenorrhea**

## **Forecast in 21 Major Markets 2016-2026**

<https://marketpublishers.com/r/E45AA7AFBBDEN.html>

Date: April 2016

Pages: 54

Price: US\$ 6,382.00 (Single User License)

ID: E45AA7AFBBDEN

### **Abstracts**

Dysmenorrhoea is a common menstrual complaint, also known as painful menstruation. It is defined as severe, painful cramping in the lower abdomen beginning shortly before or at the onset of menses. Dysmenorrhoea is often accompanied by other symptoms such as sweating, headaches, nausea, vomiting and diarrhoea. Dysmenorrhea only affects women during their fertile years.

There are two main categorisations of Dysmenorrhoea: Primary (generally associated with females less than 20 years of age with no identifiable cause) and Secondary (usually caused by presence of another pelvic condition such as endometriosis). Please note that this report mainly focuses on primary Dysmenorrhoea.

This report provides the current 12-month prevalent population for Dysmenorrhea across 21 Major Markets (USA, France, Germany, Italy, Spain, UK, Poland, Portugal, Sweden, Croatia, Switzerland, Japan, China, India, Australia, Canada, Turkey, Saudi Arabia, Egypt, Brazil and Mexico) split by 5-year age cohort. Along with the current prevalence, the report also contains a disease overview of the risk factors, disease diagnosis and prognosis along with specific variations by geography and ethnicity.

Providing a value-added level of insight from the analysis team at Black Swan, several of the main symptoms and co-morbidities of Dysmenorrhea have been quantified and presented alongside the overall prevalence figures. These sub-populations within the main disease are also included at a country level across the 10-year forecast snapshot.

Risk factors or conditions associated with developing primary Dysmenorrhea include:

BMI (low or high)

Smoking (inconclusive)

Family history of dysmenorrhea

Early age at menarche

Nulliparity

Exercise level / fitness

This report is built using data and information sourced from the proprietary Epiomic patient segmentation database. To generate accurate patient population estimates, the Epiomic database utilises a combination of several world class sources that deliver the most up to date information from patient registries, clinical trials and epidemiology studies. All of the sources used to generate the data and analysis have been identified in the report.

Reason to buy

Able to quantify patient populations in global Dysmenorrhea's market to target the development of future products, pricing strategies and launch plans.

Gain further insight into the prevalence of the subdivided types of Dysmenorrhea and identify patient segments with high potential.

Delivery of more accurate information for clinical trials in study sizing and realistic patient recruitment for various countries.

Provide a level of understanding on the impact from specific co-morbid conditions on Dysmenorrhea's prevalent population.

Identify sub-populations within Dysmenorrhea which require treatment.

Gain an understanding of the specific markets that have the largest number of Dysmenorrhea patients.

## Contents

Introduction

Cause of the Disease

Risk Factors and Prevention

Diagnosis of the Disease

Variation by Geography /Ethnicity

Disease Prognosis and Clinical Course

Key Co-Morbid Conditions / Features Associated with the Disease

Methodology for Quantifications of Patient Numbers

Top-Line Dysmenorrhea 12-Month Prevalence

Severity and Pain Frequency of Primary Dysmenorrhea

Severity of Primary Dysmenorrhea

Requirement for Medication to Treat Dysmenorrhea

Symptoms & Comorbid Conditions in Dysmenorrhea

Abbreviations used in the report

Other Black Swan Analysis Publications

Black Swan Analysis Online Patient-Based Databases

Patient-Based Offering

Online Pricing Data and Platforms

References

Appendix

LIST OF TABLES and Figures

Prevalence of Primary Dysmenorrhea, females (000s)

Prevalence of Secondary Dysmenorrhea from Endometriosis, females (000s)

Severity of Primary Dysmenorrhea patients, females (000s)

Frequency of Pain experienced by menstrual cycle, females (000s)

Duration of pain during menstrual period, females (000s)

Number of women who report taking medication for dysmenorrhea, females (000s)

Total estimated treatment days per year for dysmenorrhea, (000s)

Dysmenorrhea patients with Chronic Pelvic Pain (CPP), females (000s)

Dysmenorrhea patients with Dyspareunia, females (000s)

Dysmenorrhea patients with depression symptoms, females (000s)

Dysmenorrhea patients with headaches, females (000s)

Dysmenorrhea patients with nausea/vomiting, females (000s)

Dysmenorrhea patients with acne symptoms, females (000s)

Abbreviations and Acronyms used in the report

USA 12-month Prevalence 1° Dysmenorrhea by 5-yr age cohort, females (000s)

France 12-month Prevalence 1° Dysmenorrhea by 5-yr age cohort, females (000s)

Germany 12-month Prevalence 1° Dysmenorrhea by 5-yr age cohort, females (000s)  
Italy 12-month Prevalence 1° Dysmenorrhea by 5-yr age cohort, females (000s)  
Spain 12-month Prevalence 1° Dysmenorrhea by 5-yr age cohort, females (000s)  
UK 12-month Prevalence 1° Dysmenorrhea by 5-yr age cohort, females (000s)  
Poland 12-month Prevalence 1° Dysmenorrhea by 5-yr age cohort, females (000s)  
Portugal 12-month Prevalence 1° Dysmenorrhea by 5-yr age cohort, females (000s)  
Switzerland 12-month Prevalence 1° Dysmenorrhea by 5-yr age cohort, females (000s)  
Sweden 12-month Prevalence 1° Dysmenorrhea by 5-yr age cohort, females (000s)  
Croatia 12-month Prevalence 1° Dysmenorrhea by 5-yr age cohort, females (000s)  
Turkey 12-month Prevalence 1° Dysmenorrhea by 5-yr age cohort, females (000s)  
Japan 12-month Prevalence 1° Dysmenorrhea by 5-yr age cohort, females (000s)  
India 12-month Prevalence 1° Dysmenorrhea by 5-yr age cohort, females (000s)  
China 12-month Prevalence 1° Dysmenorrhea by 5-yr age cohort, females (000s)  
Brazil 12-month Prevalence 1° Dysmenorrhea by 5-yr age cohort, females (000s)  
Mexico 12-month Prevalence 1° Dysmenorrhea by 5-yr age cohort, females (000s)  
Australia 12-month Prevalence 1° Dysmenorrhea by 5-yr age cohort, females (000s)  
Canada 12-month Prevalence 1° Dysmenorrhea by 5-yr age cohort, females (000s)  
Egypt 12-month Prevalence 1° Dysmenorrhea by 5-yr age cohort, females (000s)  
Saudi Arabia 12-month Prevalence 1° Dysmenorrhea by 5-yr age cohort, females (000s)

## I would like to order

Product name: Epiomic Epidemiology Series: Dysmenorrhea Forecast in 21 Major Markets 2016-2026

Product link: <https://marketpublishers.com/r/E45AA7AFBBDEN.html>

Price: US\$ 6,382.00 (Single User License / Electronic Delivery)

If you want to order Corporate License or Hard Copy, please, contact our Customer Service:

[info@marketpublishers.com](mailto:info@marketpublishers.com)

## Payment

To pay by Credit Card (Visa, MasterCard, American Express, PayPal), please, click button on product page <https://marketpublishers.com/r/E45AA7AFBBDEN.html>

To pay by Wire Transfer, please, fill in your contact details in the form below:

First name:  
Last name:  
Email:  
Company:  
Address:  
City:  
Zip code:  
Country:  
Tel:  
Fax:  
Your message:

**\*\*All fields are required**

Customer signature \_\_\_\_\_

Please, note that by ordering from marketpublishers.com you are agreeing to our Terms & Conditions at <https://marketpublishers.com/docs/terms.html>

To place an order via fax simply print this form, fill in the information below and fax the completed form to +44 20 7900 3970