

Opioid-Induced Constipation - Epidemiology Forecast to 2032

https://marketpublishers.com/r/O4A610F5425EN.html

Date: January 2022

Pages: 60

Price: US\$ 3,950.00 (Single User License)

ID: O4A610F5425EN

Abstracts

This report can be delivered to the clients within 5-7 Business Days

DelveInsight's 'Opioid-Induced Constipation - Epidemiology Forecast to 2032' report delivers an in-depth understanding of the disease, historical and forecasted Opioid-Induced Constipation epidemiology in the 7MM, i.e., the United States, EU5 (Germany, Spain, Italy, France, and the United Kingdom), and Japan.

Geographies Covered

The United States

EU5 (Germany, France, Italy, Spain, and the United Kingdom)

Japan

Study Period: 2019-2032

Opioid-Induced Constipation Understanding

The DelveInsight Opioid-Induced Constipation epidemiology report gives a thorough understanding of the Opioid-Induced Constipation by including details such as disease definition, symptoms, causes, pathophysiology, and diagnosis. It also provides treatment algorithms and treatment guidelines for Opioid-Induced Constipation in the US, Europe, and Japan. The report covers the detailed information of the Opioid-Induced Constipation epidemiology scenario in seven major countries (US, EU5, and



Japan).

Opioid-Induced Constipation Epidemiology Perspective by DelveInsight

The Opioid-Induced Constipation epidemiology division provides insights about historical and current patient pool and forecasted trend for every seven major countries. The Opioid-Induced Constipation epidemiology data are studied through all possible division to give a better understanding of the Disease scenario in 7MM. The Opioid-Induced Constipation epidemiology segment covers the epidemiology data in the US, EU5 countries (Germany, Spain, Italy, France, and the UK), and Japan from 2019 to 2032. It also helps recognize the causes of current and forecasted trends by exploring numerous studies, survey reports and views of key opinion leaders.

Opioid-Induced Constipation Detailed Epidemiology Segmentation

The Opioid-Induced Constipation epidemiology covered in the report provides historical as well as forecasted Opioid-Induced Constipation epidemiology scenario in the 7MM covering the United States, EU5 countries (Germany, Spain, Italy, France, and the United Kingdom), and Japan from 2019 to 2032.

The DelveInsight Opioid-Induced Constipation report also provides the epidemiology trends observed in the 7MM during the study period, along with the assumptions undertaken. The calculated data are presented with relevant tables and graphs to give a clear view of the epidemiology at first sight.

Scope of the Report

The Opioid-Induced Constipation report covers a detailed overview explaining its causes, symptoms, classification, pathophysiology, diagnosis and treatment patterns

The Opioid-Induced Constipation Epidemiology Report and Model provide an overview of the global trends of Opioid-Induced Constipation in the seven major markets (7MM: US, France, Germany, Italy, Spain, UK, and Japan)

The report provides insight into the historical and forecasted patient pool of Opioid-Induced Constipation in seven major markets covering the United States, EU5 (Germany, Spain, France, Italy, UK), and Japan



The report helps recognize the growth opportunities in the 7MM for the patient population

The report assesses the disease risk and burden and highlights the unmet needs of Opioid-Induced Constipation

The report provides the segmentation of the Opioid-Induced Constipation epidemiology

Report Highlights

11-year Forecast of Opioid-Induced Constipation epidemiology

7MM Coverage

Prevalent and Diagnosed Cases of Opioid-Induced Constipation

Cases of Opioid-Induced Constipation by Mutation Types

Opioid-Induced Constipation Cases associated with Clinical Manifestations

KOL views

We interview, KOLs and SME's opinion through primary research to fill the data gaps and validate our secondary research. The opinion helps understand the total patient population and current treatment pattern. This will support the clients in potential upcoming novel treatment by identifying the overall scenario of the indications.

Key Questions Answered

What will be the growth opportunities in the 7MM with respect to the patient population pertaining to Opioid-Induced Constipation?

What are the key findings pertaining to the Opioid-Induced Constipation epidemiology across 7MM and which country will have the highest number of patients during the forecast period (2019-2032)?



What would be the total number of patients of Opioid-Induced Constipation across the 7MM during the forecast period (2019-2032)?

Among the EU5 countries, which country will have the highest number of patients during the forecast period (2019-2032)?

At what CAGR the patient population is expected to grow in 7MM during the forecast period (2019-2032)?

What is the disease risk, burden and unmet needs of Opioid-Induced Constipation?

What are the currently available treatments of Opioid-Induced Constipation?

Reasons to buy

The Opioid-Induced Constipation Epidemiology report will allow the user to -

Develop business strategies by understanding the trends shaping and driving the global Opioid-Induced Constipation market

Quantify patient populations in the global Opioid-Induced Constipation market to improve product design, pricing, and launch plans

Organize sales and marketing efforts by identifying the age groups and sex that present the best opportunities for Opioid-Induced Constipation therapeutics in each of the markets covered

Understand the magnitude of Opioid-Induced Constipation population by its epidemiology

The Opioid-Induced Constipation Epidemiology Model developed by DelveInsight is easy to navigate, interactive with dashboards, and epidemiology based with transparent and consistent methodologies. Moreover, the model supports data presented in the report and showcases disease trends over 11-year forecast period using reputable sources



Key Assessments

Patient Segmentation

Disease Risk & Burden

Risk of disease by the segmentation

Factors driving growth in a specific patient population



Contents

1. KEY INSIGHTS

2. EXECUTIVE SUMMARY OF OPIOID-INDUCED CONSTIPATION

3. OPIOID-INDUCED CONSTIPATION: DISEASE BACKGROUND AND OVERVIEW

- 3.1. Introduction
- 3.2. Sign and Symptoms
- 3.3. Pathophysiology
- 3.4. Risk Factors
- 3.5. Diagnosis

4. PATIENT JOURNEY

5. EPIDEMIOLOGY AND PATIENT POPULATION

- 5.1. Epidemiology Key Findings
- 5.2. Assumptions and Rationale: 7MM
- 5.3. Epidemiology Scenario: 7MM
- 5.3.1. Opioid-Induced Constipation Epidemiology Scenario in the 7MM (2019- 2032)
- 5.4. United States Epidemiology
- 5.4.1. Opioid-Induced Constipation Epidemiology Scenario in the United States (2019-2032)
- 5.5. EU-5 Country-wise Epidemiology
 - 5.5.1. Germany Epidemiology
- 5.5.1.1. Opioid-Induced Constipation Epidemiology Scenario in Germany (2019-2032)
 - 5.5.2. France Epidemiology
 - 5.5.2.1. Opioid-Induced Constipation Epidemiology Scenario in France (2019- 2032)
 - 5.5.3. Italy Epidemiology
 - 5.5.3.1. Opioid-Induced Constipation Epidemiology Scenario in Italy (2019- 2032)
 - 5.5.4. Spain Epidemiology
 - 5.5.4.1. Opioid-Induced Constipation Epidemiology Scenario in Spain (2019- 2032)
 - 5.5.5. United Kingdom Epidemiology
- 5.5.5.1. Opioid-Induced Constipation Epidemiology Scenario in the United Kingdom (2019-2032)
- 5.6. Japan Epidemiology



5.6.1. Opioid-Induced Constipation Epidemiology Scenario in Japan (2019- 2032)

6. TREATMENT ALGORITHM, CURRENT TREATMENT, AND MEDICAL PRACTICES

- 6.1. Opioid-Induced Constipation Treatment and Management
- 6.2. Opioid-Induced Constipation Treatment Algorithm
- 7. KOL VIEWS
- 8. UNMET NEEDS
- 9. APPENDIX
- 9.1. Bibliography
- 9.2. Report Methodology
- 10. DELVEINSIGHT CAPABILITIES
- 11. DISCLAIMER
- **12. ABOUT DELVEINSIGHT**

*The table of contents is not exhaustive; will be provided in the final report



List Of Tables

LIST OF TABLES

List of Table:

Table 1: Opioid-Induced Constipation Epidemiology in 7MM (2019-2032)

Table 2: Opioid-Induced Constipation Diagnosed and Treatable Cases in 7MM (2019-2032)

Table 3: Opioid-Induced Constipation Epidemiology in the United States (2019-2032)

Table 4: Opioid-Induced Constipation Diagnosed and Treatable Cases in the United States (2019-2032)

Table 5: Opioid-Induced Constipation Epidemiology in Germany (2019-2032)

Table 6: Opioid-Induced Constipation Diagnosed and Treatable Cases in Germany (2019-2032)

Table 7: Opioid-Induced Constipation Epidemiology in France (2019-2032)

Table 8: Opioid-Induced Constipation Diagnosed and Treatable Cases in France (2019-2032)

Table 9: Opioid-Induced Constipation Epidemiology in Italy (2019-2032)

Table 10: Opioid-Induced Constipation Diagnosed and Treatable Cases in Italy (2019-2032)

Table 11: Opioid-Induced Constipation Epidemiology in Spain (2019-2032)

Table 12: Opioid-Induced Constipation Diagnosed and Treatable Cases in Spain (2019-2032)

Table 13: Opioid-Induced Constipation Epidemiology in the United Kingdom (2019-2032)

Table 14: Opioid-Induced Constipation Diagnosed and Treatable Cases in the United Kingdom (2019-2032)

Table 15: Opioid-Induced Constipation Epidemiology in Japan (2019-2032)

Table 16: Opioid-Induced Constipation Diagnosed and Treatable Cases in Japan (2019-2032)



List Of Figures

LIST OF FIGURES

List of Figures

Figure 1 Opioid-Induced Constipation Epidemiology in 7MM (2019-2032)

Figure 2 Opioid-Induced Constipation Diagnosed and Treatable Cases in 7MM (2019-2032)

Figure 3 Opioid-Induced Constipation Epidemiology in the United States (2019-2032)

Figure 4 Opioid-Induced Constipation Diagnosed and Treatable Cases in the United States (2019-2032)

Figure 5 Opioid-Induced Constipation Epidemiology in Germany (2019-2032)

Figure 6 Opioid-Induced Constipation Diagnosed and Treatable Cases in Germany (2019-2032)

Figure 7 Opioid-Induced Constipation Epidemiology in France (2019-2032)

Figure 8 Opioid-Induced Constipation Diagnosed and Treatable Cases in France (2019-2032)

Figure 9 Opioid-Induced Constipation Epidemiology in Italy (2019-2032)

Figure 10 Opioid-Induced Constipation Diagnosed and Treatable Cases in Italy (2019-2032)

Figure 11 Opioid-Induced Constipation Epidemiology in Spain (2019-2032)

Figure 12 Opioid-Induced Constipation Diagnosed and Treatable Cases in Spain (2019-2032)

Figure 13 Opioid-Induced Constipation Epidemiology in the United Kingdom (2019-2032)

Figure 14 Opioid-Induced Constipation Diagnosed and Treatable Cases in the United Kingdom (2019-2032)

Figure 15 Opioid-Induced Constipation Epidemiology in Japan (2019-2032)

Figure 16 Opioid-Induced Constipation Diagnosed and Treatable Cases in Japan (2019-2032)

*The table of contents is not exhaustive; will be provided in the final report



I would like to order

Product name: Opioid-Induced Constipation - Epidemiology Forecast to 2032

Product link: https://marketpublishers.com/r/O4A610F5425EN.html

Price: US\$ 3,950.00 (Single User License / Electronic Delivery)

If you want to order Corporate License or Hard Copy, please, contact our Customer

Service:

info@marketpublishers.com

Payment

First name:

To pay by Credit Card (Visa, MasterCard, American Express, PayPal), please, click button on product page https://marketpublishers.com/r/O4A610F5425EN.html

To pay by Wire Transfer, please, fill in your contact details in the form below:

Last name:	
Email:	
Company:	
Address:	
City:	
Zip code:	
Country:	
Tel:	
Fax:	
Your message:	
	**All fields are required
	Custumer signature

Please, note that by ordering from marketpublishers.com you are agreeing to our Terms & Conditions at https://marketpublishers.com/docs/terms.html

To place an order via fax simply print this form, fill in the information below and fax the completed form to +44 20 7900 3970